



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

he [Schweigger] had based the distinction between this species [*Ch. lacertina*], considered new by him, and that which he described under the name of *Ch. serpentina*."

Therefore, if there be two forms of Snapping Turtles, their names would be *Chelydra serpentina serpentina* (Linnæus) and *Chelydra serpentina lacertina* (Schweigger).

LEONHARD STEJNEGER,

Washington, D. C.

NOTE ON HAWAIIAN GECKOS

During April, 1913, Dr. Henry A. Pilsbry secured two interesting geckos while visiting the Hawaiian Islands. As any additional information to Dr. Stejneger's account of the "Hawaiian Land Reptiles" (Proc. U. S. Nat. Mus., XXI, 1899, pp. 783-813) is desirable, I give the following records:

Peropus mutilatus (Wiegmann).

Obtained on the island of Kahoolawe, from which it has not previously been reported.

Hemidactylus garnotii Duméril and Bibron.

One secured at Moomouii, on the island of Molokai.

Lepidodactylus lugubris (Duméril and Bibron).

An old example, collected by J. K. Townsend many years ago, is interesting as likely being the first Hawaiian gecko in any museum, especially American. It is still in good preservation. Though labeled simply "Sandwich Islands," it was probably secured on Oahu. The original label also shows that it was confused with the preceding species.

H. W. FOWLER,

Philadelphia, Pa.